

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



84

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>A61K 31/505, 31/70</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 92/15309</b> <b>(43) International Publication Date:</b> 17 September 1992 (17.09.92)
<b>(21) International Application Number:</b> PCT/GB92/00389 <b>(22) International Filing Date:</b> 5 March 1992 (05.03.92) <b>(30) Priority data:</b> 9104740.7 6 March 1991 (06.03.91) GB <b>(71) Applicant (for all designated States except US):</b> THE WELL-COME FOUNDATION LIMITED [GB/GB]; Unicorn House, 160 Euston Road, London NW1 2BP (GB). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> PAINTER, George, Robert, III [US/US]; 108 Pitch Pine Lane, Chapel Hill, NC 27514 (US). FURMAN, Phillip, Allen [US/US]; 901 Bluestone Road, Durham, NC 27713 (US).		<b>(74) Agent:</b> GARRETT, M.; The Wellcome Foundation Limited, Langley Court, Beckenham, Kent BR3 3BS (GB). <b>(81) Designated States:</b> AT (European patent), AU, BE (European patent), CA, CH (European patent), CS, DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), RU, SE (European patent), US.  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ANTIVIRAL NUCLEOSIDE COMBINATION  <b>(57) Abstract</b>  Synergistic combinations of nucleoside derivatives, pharmaceutical formulations containing said combinations and use of the combinations in the treatment of retroviral infections are disclosed.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

ANTIVIRAL NUCLEOSIDE COMBINATION

The present invention relates to synergistic antiviral combinations of nucleoside derivatives, pharmaceutical formulations containing said combinations and their use in medical therapy, particularly in the treatment of virus infections, especially retrovirus infections.

Acquired immunodeficiency syndrome, (AIDS) is an immunosuppressive or immunodestructive disease that predisposes subjects to fatal opportunistic infections. Characteristically, AIDS is associated with a progressive depletion of T-cells, especially the helper-inducer subset bearing the OKT<sup>4</sup> surface marker.


Human immunodeficiency virus (HIV) has been reproducibly isolated from patients with AIDS or with the symptoms that frequently precede AIDS. HIV is cytopathic and appears to preferentially infect and destroy T-cells bearing the OKT<sup>4</sup> marker. It is now generally recognized that HIV is the etiological agent of AIDS.

Since the discovery that HIV is the etiological agent of AIDS, numerous proposals have been made for anti-HIV chemotherapeutic agents that may be effective in treating AIDS sufferers. Thus, for example, European Patent Specification 0 382 526 describes anti-HIV substituted 1,3-oxathiolanes. U.S. Patent Specification 4724232 and European Specification 0 196 185 describe 3'-azido-3'-deoxythymidine (which has the approved name zidovudine) and its use in treating AIDS.


We have now discovered that 1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine in combination with 3'-azido-3'-deoxythymidine (zidovudine) results in a surprisingly large potentiation of the anti-HIV activity of the compounds. The use of 1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine in conjunction with zidovudine produces a synergistic increase in anti-HIV activity in comparison with the anti-HIV activities of the individual compounds.

- 2 -

According to a first feature of the present invention there is provided a combination of (a) 1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine of formula (I):



or a physiologically functional derivative thereof and (b) 3'-azido-3'-deoxythymidine (zidovudine) of formula (II):



or a physiologically functional derivative thereof, components (a) and (b) of the combination being employed together such that a synergistic antiviral effect is achieved. The term "synergistic antiviral effect" is used herein to denote an antiviral effect which is greater than the predicted purely additive effects of the individual components (a) and (b) of the combination.

It should be noted that the compound of formula (I) contains two chiral centers and therefore exists in the form of two pairs of optical isomers (i.e. enantiomers) and mixtures thereof including racemic mixtures. Thus, the compound of formula (I) may be either a cis or a trans isomer or mixtures thereof. Each cis or a trans isomer or mixtures thereof. Each cis and trans isomer can exist as one of two enantiomers or mixtures thereof including racemic mixtures.

All such isomers and mixtures thereof including racemic mixtures are within the scope of the invention which also includes the tautomeric forms of the compounds of formula (I) and (II). The cis isomers of the compound of formula (I) are preferred.

By "physiologically functional derivative" is meant a pharmaceutically acceptable salt, ester or salt of an ester of the parent compound of formula (I) or (II), a pharmaceutically acceptable amide of the compound of (I), or any other compound which upon administration to the recipient is capable of providing (directly or indirectly) the parent compound or an active metabolite or residue thereof.

Preferred esters according to the invention include carboxylic acid esters in which the non-carbonyl moiety of the ester grouping is selected from straight or branched chain alkyl e.g. n-propyl, t-butyl, n-butyl, alkoxyalkyl (e.g. methoxymethyl), aralkyl (e.g. benzyl), aryloxyalkyl (e.g. phenoxymethyl), and aryl (e.g. phenyl); sulfonate esters such as alkyl- or aralkylsulfonyl (e.g. methanesulfonyl); amino acid esters (e.g. L-valyl or L-isoleucyl); dicarboxylic acid esters (e.g., hemisuccinate); and mono-, di- or triphosphate esters. The phosphate esters may be further esterified by, for example, a  $C_{1-20}$  alcohol or reactive derivative thereof, or by a 2,3-di( $C_{6-24}$ )acyl glycerol.

Any alkyl moiety present in such esters advantageously contains 1 to 18 carbon atoms, particularly 1 to 4 carbon atoms. Any aryl moiety present in such esters advantageously comprises a phenyl group optionally substituted, e.g. by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy or nitro.

The above-mentioned pharmaceutically acceptable amides of the compound of formula (I) include those derivatives wherein the cytosine amino group is present in the form of an amide, eg.  $NHCOR$  wherein R is  $C_{1-6}$  alkyl or aryl (eg. phenyl optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, nitro or hydroxyl).

Examples of pharmaceutically acceptable salts include base salts, e.g. derived from an appropriate base, such as alkali metal (e.g. sodium), alkaline earth metal (e.g. magnesium) salts, ammonium and  $NX_4^+$  (wherein X is  $C_{1-4}$  alkyl). Pharmaceutically acceptable acid addition salts include salts of organic carboxylic acids such as acetic, lactic, tartaric, malic, isethionic, lactobionic and succinic acids; organic sulfonic acids such as methanesulfonic, ethanesulfonic, benzenesulfonic and p-toluenesulfonic acids and inorganic acids such as hydrochloric, sulfuric, phosphoric and sulfamic acids.

Examples of viral infections and associated clinical conditions which may be treated or prevented in accordance with the invention, include human retroviral infections such as human immunodeficiency virus (HIV), e.g. HIV-1 or HIV-2, and human T-cell lymphotropic virus (HTLV), e.g. HTLV-I or HTLV-II infections. The combinations of the present invention are especially useful for the treatment of AIDS and related clinical conditions such as AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), AIDS-related neurological conditions, such as multiple sclerosis or tropical paraparesis, anti-HIV antibody-positive and HIV-positive conditions such as thrombocytopenic purpura. The combinations of the present invention may also be used in the treatment of psoriasis. The combinations of the present invention have been found to be particularly applicable to the treatment of asymptomatic infections or diseases caused by or associated with human retroviruses.

According to a second feature of the invention there are provided combinations as hereinbefore described, for use in medical therapy, particularly for the treatment or prophylaxis of any of the aforementioned viral infections or conditions, especially HIV infections including AIDS.

The present invention further includes a process for preparing the combinations hereinbefore described, which comprises bringing into

association components (a) and (b) of the combination in a medicament to provide a synergistic antiviral effect.

In a further aspect of the present invention, there is provided the use of a combination of the present invention in the manufacture of a medicament for the treatment of any of the aforementioned viral infections or conditions.

The present invention further provides a method for the treatment or prophylaxis of viral infections (especially HIV infections) in a mammal (including a human) which comprises administering to said mammal an effective amount of a combination as hereinbefore described. It will be appreciated that in accordance with the present invention, components (a) and (b) of the combination may be administered simultaneously or sequentially. In the latter case, however, the components are administered within a sufficiently short interval to ensure that a synergistic antiviral effect is achieved.

The present invention also provides a method of potentiating in a mammal (including a human) having a viral infection, the antiviral activities of components (a) and (b) of the combination, which comprises administering to said mammal an effective synergistic amount of component (a) simultaneously with, previous to, or subsequent to the administration of component (b).

An advantage of the combination of the present invention is that it enables attainment of an improved antiviral efficacy at a particular dose of one of the antiviral components (compared with the component used alone) thereby improving the therapeutic index of the component. Thus, for example, the combination may be used to treat conditions which would otherwise require relatively large doses of the antiviral component at which toxicity problems may occur. The smaller doses of the combination may provide increased convenience to the recipient and increased compliance.



The combinations of the present invention may be administered to a mammal in a conventional manner. As indicated above, components (a) and (b) may be administered simultaneously (e.g., in a unitary pharmaceutical formulation) or separately (e.g., in separate pharmaceutical formulations). In general, the combinations may be administered by the topical, oral, rectal or parenteral (e.g., intravenous, subcutaneous or intramuscular) routes. It will be appreciated that the route may vary with, for example, the severity of the condition to be treated and the identity of the recipient.

It will be appreciated that, while there will usually be an optimum ratio of the components to ensure maximum potentiation, even a vanishingly small quantity of one component will suffice to potentiate the effect of the other to some degree, and so any ratio of two potentiating components will still possess the required synergistic effect. However, greatest synergy is generally observed when the two components are present in particular ratios.

Thus the optimum molar ratios of zidovudine to 1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine, or their respective physiologically functional derivatives, for use according to this invention are from 1:1 to 1:600, preferably from 1:10 to 1:250, and most preferably 1:25.

Hereafter the components of the combination may be referred to as "active ingredients".

The dose of the combination will depend on the condition being treated and other clinical factors such as the weight and condition of the recipient and the route of administration of the components of the combinations. Examples of dose ranges and component ratios are as follows:

In general a suitable dose of a combination of the present invention based on the total weight of components (a) and (b) will be in the

range of 3 to 120 mg per kilogram body weight of the recipient per day, preferably in the range of 6 to 90 mg per kilogram body weight per day and most preferably in the range 15 to 60 mg per kilogram body weight per day. The desired dose is preferably presented as two, three, four, five, six or more sub-doses administered at appropriate intervals throughout the day. These sub-doses may be administered in unit dose forms, for example, containing 10 to 1500 mg, preferably 20 to 1000 mg, and most preferably 50 to 700 mg of active ingredients per unit dose form.

While it is possible for the active ingredients to be administered alone it is preferable to present them as pharmaceutical formulations. Pharmaceutical formulations of the present invention comprise a combination according to the invention together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. When the individual components of the combination are administered separately they are generally each presented as a pharmaceutical formulation. The references hereinafter to formulations refer unless otherwise stated to formulations containing either the combination or a component thereof. Formulations include those suitable for oral, rectal, nasal, topical (including transdermal, buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. Such methods include the step of bringing into association the active ingredients with the carrier which constitutes one or more accessory ingredients.

In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredients; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as a powder or granules, optionally mixed with a binder (e.g. povidone, gelatin, hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (e.g. sodium starch glycollate, cross-linked povidone, cross-linked sodium carboxymethyl cellulose) surface-active or dispersing agent. Molded tablets may be made by molding a mixture of the powdered compound moistened with an inert liquid diluent in a suitable machine. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredients therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile. Tablets may optionally be provided with an enteric coating, to provide release in parts of the gut other than the stomach.

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredients in a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier. Formulations for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or a salicylate.

Topical administration may also be by means of a transdermal iontophoretic device.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and nonaqueous isotonic sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents; and liposomes or other microparticulate systems which are designed to target the compound to blood components or one or more organs. The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.


Preferred unit dosage formulations are those containing a daily dose or daily subdose of the active ingredients, as hereinbefore recited, or an appropriate fraction thereof.

It should be understood that in addition to the ingredients particularly mentioned above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example, those suitable for oral administration may include such further agents as sweeteners, thickeners and flavoring agents.

The compounds of the combination of the present invention may be prepared in conventional manner. Zidovudine can be prepared, for example, as described in U.S. Patent 4724232, incorporated herein by reference. Zidovudine can also be obtained from Aldrich Chemical Co., Milwaukee, WI 53233, USA.


1-(2-(Hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine may be prepared for example by:

- a) reacting optionally protected 5-fluorocytosine with a 1,3-oxathiolane of formula (IIIA):




wherein  $R_1$  is hydrogen or a hydroxy protecting group and L is a leaving group; or

- b) reacting a compound of formula (IIIB):



(wherein  $R_1$  is as defined above and  $R_1^a$  is an amino protecting group) with a fluorinating agent serving to introduce a fluorine atom in the 5-position of the cytosine ring; or

c) reacting a compound of formula (IIIC):



(wherein  $R_1$  is as defined above) with an agent serving to convert the oxo group in the 4-position of the uracil ring to an amino group; any remaining protecting groups being removed to produce the desired product.

With the regard to process a), the hydroxy protecting group includes protecting groups such as acyl (e.g. acetyl), arylacyl (e.g. benzoyl or substituted benzoyl), trityl or monomethoxytrityl, benzyl or substituted benzyl, trialkylsilyl (e.g. dimethyl-*t*-butylsilyl) or diphenylmethylsilyl. The 5-fluorocytosine compound may be optionally protected with silyl, e.g. trimethyl silyl groups. Such groups may be removed in conventional manner. The leaving group L is a leaving group typical of those known in the art of nucleoside chemistry e.g. halogen such as chlorine or bromine, alkoxy such as methoxy or ethoxy or acyl such as acetyl or benzoyl.

The reaction in process a) may be affected in an organic solvent (e.g. 1,2-dichloroethane or acetonitrile) in the presence of a Lewis acid such as stannic chloride or trimethylsilyl triflate.

Compounds of formula IIIA may be obtained from a suitably protected 2-hydroxyacetaldehyde of formula (IV):



wherein  $R_1$  is defined above, as described in Can. J. Research, 8, 129 (1933) and European Patent Specification 0 382 526. Reaction of compounds of formula IV with a mercaptoacetal  $HSCH_2CH(OR)_2$  wherein R is  $C_{1-4}$  alkoxy such as  $HSCH_2CH(OC_2H_5)_2$ , known in the art (Chem. Ber. 85, 924-932 (1952)), yields compounds of formula IIIA wherein L is OR (alkoxy) e.g. methoxy or ethoxy. Alternatively, compounds of formula IIIA, wherein L is alkoxy, may be converted to compounds of formula IIIA wherein L is halogen or acyl by methods known in the art of carbohydrate chemistry.

Compounds of formula IV may be prepared from 1,2-O-isopropylidene glycerol by introduction of  $R_1$  (e.g. trisubstituted silyl, benzyl or trityl) and removal of the isopropylidene group with mild acid (e.g. aqueous formic or acetic acid) or zinc bromide in acetonitrile, followed by oxidation of the alcohol group with aqueous periodate.

With regard to process b), the 5-fluoro substituent may be introduced by methods known in the art (M.J. Robins, et al., in Nucleic Acid Chemistry. Part 2, L.B. Townsend and R.S. Tipson, editors, J. Wiley and Sons, New York, 895-900 (1978) and references therein; R. Duschinsky in Nucleic Acid Chemistry, Part 1, L.B. Townsend and R.S. Tipson, editors, J. Wiley and Sons, New York, 43-46 (1978) and references therein). The fluorinating agent may be, for example, trimethylhypofluorite in fluorotrichloromethane.

With regard to process c), the compound of formula (IIIC) is advantageously treated with 1,2,4-triazole, advantageously together with 4-chlorophenyl dichlorophosphate, to form the corresponding 4-(1,2,4-triazolyl) compound which is then converted to the desired 4-amino (cytidine) compound by reaction with for example methanol.

The starting materials of formula IIIB and IIIC may be prepared for example by reaction of an appropriate (optionally protected) base with a compound of formula IIIA in an analogous manner to that described in process a). 5-Fluorouracil and 5-fluorocytosine are commercially available from Aldrich Chemical Co., Milwaukee, WI 53233, USA.

Separation of the ( $\pm$ )-cis and ( $\pm$ )-trans isomers of formula (I) for example in a protected form, may be accomplished by chromatography on silica gel with mixtures of organic solvents such as ethyl acetate/methanol, ethyl acetate/hexane or dichloromethane/methanol. Any protecting group may then be removed using the appropriate reagent for each group.

Esters of the component compounds of formulas I and II may be prepared in conventional manner by reaction with an appropriate esterifying agent such as an acid halide or anhydride. The compounds of formulas I and II or esters thereof may be converted into pharmaceutically acceptable salts thereof by treatment with an appropriate base. An ester or salt of the component compounds may be converted into the parent compound by hydrolysis.

Pharmaceutically acceptable amides of the compound of formula (I) may be prepared, for example by reaction with an appropriate acylating agent, for example, an acid halide or anhydride serving to acylate the 5'-OH and 4-NH<sub>2</sub> groups. The acyl group may then be removed selectively from one or other of the 5'-OH and 4-NH<sub>2</sub> groups. For example, treatment of the diacylated compound under acidic conditions, eg. a Lewis acid such as zinc bromide in methanol, removes the 4N-acyl group to yield the corresponding 5'-OH ester, while treatment of the diacylated compound under alkaline conditions, eg. with sodium methoxide removes the 5'-OH acyl group to yield the corresponding 4N-amide. The acyl groups can also be removed selectively by treatment with commercially available esterase or lipase enzymes, eg. pig liver esterase or pancreatic lipase, or by treatment in accordance with methods described in U.S. Patent Specification No. 5071983. The



compound of formula (I) may be converted into a pharmaceutically acceptable salt thereof in a conventional manner, for example, by treatment with an appropriate base.

The following examples are intended for illustration only and are not intended to limit the scope of the invention in any way. "Active ingredient" denotes a mixture of the components zidovudine and cis-1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine in the molar ratio of 1:25.

Example 1: Tablet Formulation

The following formulations A, B and C are prepared by wet granulation of the ingredients with a solution of povidone, followed by addition of magnesium stearate and compression.

Formulation A

	<u>mg/tablet</u>
Active Ingredient	250
Lactose B.P.	210
Povidone B.P.	15
Sodium Starch Glycollate	20
Magnesium Stearate	<u>5</u>
	500

Formulation B

	<u>mg/tablet</u>
Active Ingredient	250
Lactose B.P.	150
Avicel PH 101	60
Povidone B.P.	15
Sodium Starch Glycollate	20
Magnesium Stearate	<u>5</u>
	500

Formulation C

	<u>mg/tablet</u>
Active Ingredient	250
Lactose B.P.	200
Starch	50
Povidone	5
Magnesium Stearate	<u>4</u>
	359

The following formulations, D and E, are prepared by direct compression of the admixed ingredients. The lactose in formulation E is of the direct compression type (Dairy Crest - "Zeparox").

Formulation D

	<u>mg/tablet</u>
Active Ingredient	250
Pregelatinized Starch NF15	<u>150</u>
	400

Formulation E

	<u>mg/tablet</u>
Active Ingredient	250
Lactose B.P.	150
Avicel	<u>100</u>
	500

Formulation F (Controlled Release Formulation)

The formulation is prepared by wet granulation of the ingredients with a solution of povidone followed by the addition of magnesium stearate and compression.

	<u>mg/tablet</u>
Active Ingredient	500
Hydroxypropylmethylcellulose (Methocel K4M Premium)	112
Lactose B.P.	53
Povidone B.P.	28
Magnesium Stearate	<u>7</u>
	700

Drug release takes place over a period of about 6-8 hours and is complete after 12 hours.

Example 2: Capsule FormulationsFormulation A

A capsule formulation is prepared by admixing the ingredients of formulation D in Example 1 above and filling into a two-part hard gelatin capsule. Formulation B (infra) is prepared in a similar manner.

Formulation B

	<u>mg/capsule</u>
Active Ingredient	250
Lactose B.P.	143
Sodium Starch Glycollate	25
Magnesium Stearate	<u>2</u>
	420

Formulation C

	<u>mg/capsule</u>
Active Ingredient	250
Macrogel 4000 B.P.	<u>350</u>
	600

Capsules of formulation C are prepared by melting the Macrogel 4000 B.P., dispersing the active ingredient in the melt and filling the melt into a two-part hard gelatin capsule.

Formulation D

	<u>mg/capsule</u>
Active Ingredient	250
Lecithin	100
Arachis Oil	<u>100</u>
	450

Capsules of formulation D are prepared by dispersing the active ingredient in the lecithin and arachis oil and filling the dispersion into soft, elastic gelatin capsules.

Formulation E (Controlled Release Capsule)

The following controlled release capsule formulation is prepared by extruding ingredients a, b, and c using an extruder, followed by spheronization of the extrudate and drying. The dried pellets are then coated with release-controlling membrane (d) and filled into a two-piece, hard gelatin capsule.

	<u>mg/capsule</u>
(a) Active Ingredient	250
(b) Microcrystalline Cellulose	125
(c) Lactose B.P.	125
(d) Ethyl Cellulose	<u>13</u>
	513

Example 3: Injectable FormulationFormulation A

	<u>mg</u>
Active Ingredient	200
Hydrochloric Acid Solution 0.1 M <u>or</u>	
Sodium Hydroxide Solution 0.1 M q.s. to pH	4.0 to 7.0
Sterile water q.s. to	10 ml

The active ingredient is dissolved in most of the water (35°-40°C) and the pH adjusted to between 4.0 and 7.0 with the hydrochloric acid or the sodium hydroxide as appropriate. The batch is then made up to volume with the water and filtered through a sterile micropore filter into a sterile 10 ml amber glass vial (type 1) and sealed with sterile closures and overseals.

Formulation B

Active Ingredient	125 mg
Sterile, Pyrogen-free, pH 7 Phosphate	
Buffer, q. s. to	25 ml

Example 4: Intramuscular injection

Active Ingredient	200 mg
Benzyl Alcohol	0.10 g
Glycofurol 75	1.45 g
Water for injection q.s. to	3.00 ml

The active ingredient is dissolved in the glycofurol. The benzyl alcohol is then added and dissolved, and water added to 3 ml. The mixture is then filtered through a sterile micropore filter and sealed in sterile 3 ml amber glass vials (type 1).

Example 5: Syrup

Active Ingredient	250 mg
Sorbitol Solution	1.50 g
Glycerol	2.00 g
Sodium Benzoate	0.005 g
Flavor, Peach 17.42.3169	0.0125 ml
Purified Water q.s. to	5.00 ml

The active ingredient is dissolved in a mixture of the glycerol and most of the purified water. An aqueous solution of the sodium benzoate is then added to the solution, followed by addition of the sorbital solution and finally the flavor. The volume is made up with purified water and mixed well.

Example 6: Suppositorymg/capsule suppository

Active Ingredient	250
Hard Fat, B.P. (Witepsol H15 - Dynamit Nobel)	<u>1770</u>
	2020

One-fifth of the Witepsol H15 is melted in a steam-jacketed pan at 45°C maximum. The active ingredient is sifted through a 200µm sieve and added to the molten base with mixing, using a Silverson fitted with a cutting head, until a smooth dispersion is achieved. Maintaining the mixture at 45°C, the remaining Witepsol H15 is added to the suspension and stirred to ensure a homogenous mix. The entire suspension is passed through a 250µm stainless steel screen and, with continuous stirring, is allowed to cool to 40°C. At a temperature of 38°C to 40°C, 2.02 g of the mixture is filled into suitable, 2 ml plastic molds. The suppositories are allowed to cool to room temperature.

Example 7: Pessariesmg/pessary

Active Ingredient	250
Anhydrate Dextrose	380
Potato Starch	363
Magnesium Stearate	<u>7</u>
	1000

The above ingredients are mixed directly and pessaries prepared by direct compression of the resulting mixture.



Example 8: Preparation of 1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine

Method A: (±)-cis and (±)-trans 2-benzoyloxymethyl-5-(N<sub>4</sub>-acetylcytosin-1-yl)-1,3-oxathiolane are prepared and separated to the (±)-cis and (±)-trans isomers as described in European Patent (EP) Specification 0 382 526. The (±)-cis isomer is fluorinated with trifluoromethyl hypofluorite in fluorotrichloromethane (CCl<sub>3</sub>F) and chloroform at -78°C, according to the method of Robins, et al. Nucleic Acid Chemistry, Part 2, 895-900 (1978). The N<sub>4</sub>-acetyl and 2-benzoyl groups are removed with dimethylamine in ethanol, and the product, (±)-cis-1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine, is isolated.

Method B: (±)-cis and (±)-trans 2-benzoyloxymethyl-5-(uracil-1-yl)-1,3-oxathiolane are prepared as described in EP 0 382 526). After deprotection of the 2-hydroxyl group with saturated methanolic ammonia, the isomers are separated on silica gel using EtOAc/MeOH as eluant (EP 0 382 526). The (±)-cis isomer is reacted with acetic anhydride in pyridine at room temperature to give the 2-acetate. Solvent is removed in vacuo at <30°C. The 2-acetate is then dissolved in CHCl<sub>3</sub> and washed with aqueous bicarbonate. The separated organic layer is dried, and CHCl<sub>3</sub> is evaporated in vacuo. (±)-cis-2-Acetyloxymethyl-5-(uracil-1-yl)-1,3-oxathiolane is fluorinated as described above (Method A) by the method of Robins et al. Conversion of the 5-F-uracil base to the 5-F-cytosine base is carried out by preparation of the 4-(1,2,4-triazol-1-yl) derivative according to the methods of C.B. Reese, J. Chem. Soc., Perkins I, 1171 (1984) and W.L. Sung, Nucleic Acids Res., 9, 6139 (1981), using 1,2,4-triazole and 2 equivalents of 4-chlorophenyldichlorophosphate in dry pyridine at ambient temperature. This conversion is followed by reaction with methanol previously saturated with ammonia at 0°C, and the 2-acetate is hydrolyzed to give (±)-cis-1-(2-hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine.

### Antiviral Activity

Combinations according to the invention were tested for anti-HIV activity in an HIV-infected MT4 cell assay as described in Averett, D.R., J. Virol. Methods, 23, 263-276 (1989). The cells were exposed to HIV for one hour prior to addition of antiviral component(s). Components were tested in serial 2.5-fold dilutions. After five days of incubation at 37°C, the cell number was determined. Inhibition of HIV-induced cytopathic effect was calculated, and synergism was determined by FIC plots as described by Elion, Singer, and Hitchings, J. Biol. Chem. 208, 477 (1954).

The fractional inhibitor concentrations (FIC) of zidovudine and cis-1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine were calculated according to the method of Elion et al., supra (Table 1).

These values can be plotted on a graph from which it can be determined that the combination of cis-1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine and zidovudine is strongly synergistic.

Table 1

## Calculation of Fractional Inhibitor

## Concentration (FIC)

## 70% Inhibition

Zidovudine ( $\mu$ M)	Compound 1 <sup>*</sup> ( $\mu$ M)	FIC Zidovudine	FIC Compound 1
0.004	2.5	0.018	0.48
0.01	2.0	0.045	0.38
0.0256	1.6	0.12	0.31
0.06	1.4	0.27	0.26
0.22	-		
-	5.2		

\* Compound 1 is cis-1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine.

CLAIMS


1. A pharmaceutical combination comprising the components:
  - (a) 1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine or a physiologically functional derivative thereof, and
  - (b) 3'-azido-3'-deoxythymidine or a physiologically functional derivative thereof,components (a) and (b) of the combination being employed in a ratio whereby a synergistic anti-HIV effect is achieved.
2. A pharmaceutical combination as claimed in claim 1 wherein component (a) is cis-1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine and component (b) is 3'-azido-3'-deoxythymidine.
3. A pharmaceutical combination as claimed in claim 1 or claim 2 wherein the components are employed in a molar ratio in the range from 600:1 to 1:1 of component (a) to component (b).
4. A pharmaceutical combination as claimed in claim 3 wherein the components are employed in a molar ratio in the range from 250:1 to 10:1 of component (a) to component (b).
5. A pharmaceutical combination as claimed in the preceding claims for use in human therapy.
6. A pharmaceutical combination as claimed in claim 5 for use in the treatment or prophylaxis of human immunodeficiency virus (HIV) infections.
7. Use of a pharmaceutical combination as claimed in any of claims 1 to 4 in the manufacture of a medicament for use in the treatment or prophylaxis of HIV infections.

8. A process for the preparation of a pharmaceutical combination as claimed in any of claims 1 to 4 which comprises bringing into association said components (a) and (b) of the combination to provide a synergistic antiviral effect.
9. A pharmaceutical formulation comprising a pharmaceutical combination as claimed in any of claims 1 to 4 together with one or more pharmaceutically acceptable carriers or excipients.
10. A pharmaceutical formulation as claimed in claim 9 in the form of a tablet or capsule.
11. A method for the treatment or prophylaxis of a HIV infection in a human which comprises administering to said human an effective amount of a combination as claimed in any of claims 1 to 6.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 92/00389

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.C1.5                      A 61 K 31/505                      A 61 K 31/70		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.C1.5	A 61 K 31/00                      C 07 D 411/00	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>*</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
Y	US,A,4724232 (J.L. RIDEOUT et al.) 9 February 1988, see abstract; claims 1-3; column 1, line 44 - column 2, line 42; examples 7,8; column 4, lines 7-18 (cited in the application) -----	1,2,5- 10
Y	EP,A,0382526 (IAF BIOCHEM INTERNATIONAL INC.) 16 August 1990, see abstract; claims 1-5,10-13,28; page 9, lines 6-25; page 9, line 50 - page 10, line 15; page 11, lines 2-20 (cited in the application) -----	1,2,5- 10
<p><sup>*</sup> Special categories of cited documents: <sup>10</sup></p> <p><sup>"A"</sup> document defining the general state of the art which is not considered to be of particular relevance</p> <p><sup>"E"</sup> earlier document but published on or after the international filing date</p> <p><sup>"L"</sup> document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p><sup>"O"</sup> document referring to an oral disclosure, use, exhibition or other means</p> <p><sup>"P"</sup> document published prior to the international filing date but later than the priority date claimed</p> <p><sup>"T"</sup> later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p><sup>"X"</sup> document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p><sup>"Y"</sup> document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p><sup>"A"</sup> document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
12-05-1992	3 JUN 1992	
International Searching Authority	Signature of Authorized Officer	
EUR PEAN PATENT OFFICE	Mme N. KUIPER 	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE:

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 11, because they relate to subject matter not required to be searched by this Authority, namely:

See PCT Rule 39.1 (iv)  
Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.

2. ☐ Claim numbers \_\_\_\_\_, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers \_\_\_\_\_, because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 8.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING:

This international Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the international Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.  
☐ No protest accompanied the payment of additional search fees.